

Fits, faints and funny turns— Advice Sheet

Advice for parents and carers



GIG
CYMRU
NHS
WALES

Bwrdd Iechyd Prifysgol
Aneurin Bevan
University Health Board



There are many reasons for childhood “funny turns”. Your medical practitioner will have given you a likely diagnosis and this leaflet contains links for further information on some of the commonest causes of funny turns in childhood. This leaflet is designed to alert you should your child’s symptoms change and advice as to where to get help.

It helps to have video of your child’s episodes to take to any medical appointments.

When should you worry?



RED

If your child has any of the following at anytime:

- If they become unconscious and do not recover after 5 minutes
- If they have a seizure lasting more than 5 minutes
- If they have >3 episodes in one day
- Becomes confused or unaware of their surroundings (for longer than 30 mins)
- Develops difficulty speaking or understanding what you are saying
- Develops weakness in their arms/ legs or starts losing their balance
- Develops problems with their eyesight
- Collapses during exercise/ strenuous activity

You need urgent help.

Go to the nearest
Hospital Emergency
(A&E) Department or
phone 999



AMBER

If your child has any of the following at anytime:

- If they have more than one convulsive seizure in a week (Convulsive = unconscious with arms and legs stiff, sometimes shaking)
- Loss of consciousness but has quick recovery (less than 5 minutes)
- Change in their behaviour or excessive tiredness
- Develops a persistent headache that doesn't go away (despite painkillers such as paracetamol or ibuprofen)
- Develops a headache that wakes them from sleep

**You need to contact a
doctor or nurse today.**

Please ring your GP
surgery or contact
NHS 111 - dial 111 or
for children aged 5
years and above visit
111.wales.nhs.uk



GREEN

Your child may have been diagnosed with one of the below conditions that requires no medical treatment.

Here are links for further information.

Breath holding attacks: www.nhs.uk/conditions/breath-holding-spells-in-children/

Simple faint: www.nhs.uk/conditions/fainting/

Reflex anoxic seizures: www.hearhythmalliance.org/stars/uk/reflex-anoxic-seizures-ras

Night terrors <https://www.nhs.uk/conditions/night-terrors/>

Sleep myoclonus (Brief jerks of one or more limbs. Occurs only in sleep. No treatment needed)

Self Care

Continue providing your
child's care at home.

If you are still
concerned about your
child, contact NHS 111
– dial 111 or for children
aged 5 years and
above visit

111.wales.nhs.uk

www.abbHealthierTogether.cymru.nhs.uk