Lymphadenopathy advice sheet

Advice intended for parents/ carers taking their child home after seeing a healthcare professional







How is your child?



How can I help my child?

Use painkillers such as ibuprofen and paracetamol to keep your child comfortable - please read and follow the instructions on the medicine container

www.abbHealthierTogether.cymru.nhs.uk

Information based on National Health Care Guidelines and adapted from National Healthier Together Programme

Lymphadenopathy advice sheet

Advice intended for parents/ carers taking their child home after seeing a healthcare professional







Why does your child have enlarged lymph nodes?

- It is normal for lymph nodes in your child's neck to be enlarged when they have an infection such as a sore throat. This is your child's normal response to fighting common infections. Antibiotics are not normally required.
- Children with severe eczema commonly have enlarged lymph nodes. This will improve with treatment of your child's eczema.

What should you look out for?

- Occasionally, enlarged lymph nodes can become infected. If the lymph node is painful, red and hot, your child will need to see a healthcare professional because they may need treatment with antibiotics.
- If your child has been prescribed antibiotics for an infection of their lymph node and still has a fever after 2 days, they will need to be seen again.

How long will it take for my child to get better?

Your child should start getting better within a couple of days but their lymph nodes may take 2-4 weeks to improve. Small lymph nodes may persist for years.



Painless enlarged lymph nodes on both sides of the neck (bilateral) associated with a sore throat - likely to improve without treatment



Painful, hot swelling on left side of neck caused by an infected lymph node – requires treatment with antibiotics

CS51313

www.abbHealthierTogether.cymru.nhs.uk

Information based on National Health Care Guidelines and adapted from National Healthier Together Programme